



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Since $x=h$ when $t=0$, $C=\frac{4K\sqrt{h}}{ab\sqrt{2g}}$.

$$\therefore t=T=\frac{4K}{ab\sqrt{2g}}(\sqrt{h}-\sqrt{x}). \quad \therefore x=\left(\frac{4K\sqrt{h}-Tab\sqrt{2g}}{4K}\right)^2.$$

AVERAGE AND PROBABILITY.

98. Proposed by REV. PREBENDARY WHITWORTH, A. M.

A has £ m and B has £ n . They play for points until one of them has lost all his money. If α and β be the respective chances that A and B win any point, the expectation of the number of points played will be

$$\frac{n\alpha^n(\alpha^m-\beta^m)-m\beta^m(\alpha^n+\beta^n)}{(\alpha-\beta)(\alpha^{m+n}-\beta^{m+n})}.$$

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let A_m =A's chance of winning, B_n =B's chance of winning.

Then nA_m, mB_n =A's and B's expectation, respectively.

\therefore Expectation of number of points played= E .

Then $E=(nA_m-mB_n)/(\alpha-\beta)$.

Let A_x =A's chance when he has x pounds and B has $m+n-x$ pounds.

$$\therefore A_x=\frac{\beta}{\alpha+\beta}A_{x-1}+\frac{\alpha}{\alpha+\beta}A_{x+1}.$$

$A_x-A_{x-1}=(\alpha/\beta)(A_{x+1}-A_x)$. Giving x successive values from 1 to x we get $A_1-A_0=(\alpha/\beta)(A_2-A_1)$, $A_2-A_1=(\alpha/\beta)(A_3-A_2)$, etc.

By continued multiplication we get $A_1-A_0=(\alpha/\beta)^{x-1}(A_x-A_{x-1})$ or $A_x-A_{x-1}=(\beta/\alpha)^{x-1}(A_1-A_0)$.

Give x successive values from 1 to x and add

$$A_x-A_0=(A_1-A_0)[1+\beta/\alpha+(\beta/\alpha)^2+\dots+(\beta/\alpha)^{x-1}].$$

But $A_0=0$. $\therefore A_x=A_1[1-(\beta/\alpha)^x]/[1-(\beta/\alpha)]$.

$$A_{m+n}=1. \quad \therefore 1=A_1[\alpha^{m+n}-\beta^{m+n}]/[\alpha^{m+n-1}(\alpha-\beta)].$$

$$\therefore A_1=[\alpha^{m+n-1}(\alpha-\beta)]/(\alpha^{m+n}-\beta^{m+n}).$$

$$\therefore A_x=[\alpha^{m+n-1}(\alpha^x-\beta^x)]/[\alpha^{x-1}(\alpha^{m+n}-\beta^{m+n})].$$

$$\therefore A_m=[\alpha^n(\alpha^m-\beta^m)]/(\alpha^{m+n}-\beta^{m+n}).$$

$$\text{Similarly, } B_n=[\beta^m(\alpha^n-\beta^n)]/(\alpha^{m+n}-\beta^{m+n}).$$

$$\therefore E=\frac{n\alpha^n(\alpha^m-\beta^m)-m\beta^m(\alpha^n-\beta^n)}{(\alpha-\beta)(\alpha^{m+n}-\beta^{m+n})}.$$

99. Proposed by E. B. SEITZ.

A point is taken at random in the surface of a given circle, and from it a line equal in length to the radius is drawn, so as to lie wholly in the surface of the circle. Find the chance that the line intersects in a given diameter. [No. 135, *The Analyst*.]